

UNDERAGE DRINKING IN NEW YORK


The Facts

Tragic health, social, and economic problems result from the use of alcohol by youth. Underage drinking is a causal factor in a host of serious problems, including homicide, suicide, traumatic injury, drowning, burns, violent and property crime, high-risk sex, fetal alcohol syndrome, alcohol poisoning, and the need for treatment of alcohol abuse and dependence.

Problems and Costs Associated with Underage Drinking in New York

In 2013, underage drinking cost the citizens of New York \$3.1 billion. These costs include medical care, work loss, and pain and suffering associated with the multiple problems resulting from the use of alcohol by youth.¹ This translates to \$1,706 per year for each youth in the state or \$3.35 per drink consumed underage. Excluding pain and suffering from these costs, tangible costs of underage drinking including medical care, criminal justice, property damage, and loss of work in New York totaled \$1.01 billion each year or \$1.18 per drink. In contrast, a drink in New York retails for \$1.29.

**Costs of Underage Drinking
New York, 2013 \$**



**Costs of Underage Drinking by Problem,
New York, 2013 \$**

Problem	Total Costs (in millions)
Youth violence	\$1,900.8
Youth traffic crashes	\$331.7
High-risk sex, Ages 14–20 years	\$194.1
Property and public order crime	\$10.8
Youth injury	\$155.3
Poisonings and psychoses	\$52.8
Fetal alcohol syndrome among mothers aged 15–20 years	\$79.9
Youth alcohol treatment	\$179.2
Total	\$3,117.4 (e.g. \$3.1 B)

Youth violence (homicide, suicide, aggravated assault) and traffic crashes attributable to alcohol use by underage youth in New York represent the largest costs for the state. However, a host of other problems contribute substantially to the overall cost. Among teen mothers, fetal alcohol syndrome alone costs New York \$80 million.

In 2012, 3,462 youth aged 12 to 20 years were admitted for alcohol treatment in New York, accounting for 3% of all treatment admissions for alcohol abuse in the state.² Young people who begin drinking before age 15 are four times more likely to develop alcohol dependence and are two and a half times more likely to become

abusers of alcohol than those who begin drinking at age 21.³ We did not cost these adult problems.

Alcohol Consumption by Youth in New York

Underage drinking is widespread in New York. Approximately 640,000 underage customers drink each year in New York. In 2013, New York students in grades 9 to 12 reported the following:⁴

- 63.5% had at least one drink of alcohol on one or more days during their life.⁵
- 19.0% had their first drink of alcohol, other than a few sips, before age 13.
- 32.5% had at least one drink of alcohol on one or more occasions in the past 30 days.
- 18.4% had five or more drinks of alcohol in a row (binge drinking) in the past 30 days.

In 2012, underage customers consumed 10.2% of all alcohol sold in New York, totaling \$1.2 billion in sales (in 2013 dollars). These sales provided profits of \$587 million to the alcohol industry.¹ Ranking states based on the percentage of alcohol consumed underage, with 1 the highest, New York ranked number 16. This percentage is affected by both adult and youth drinking levels.

Annual sales of alcohol consumed by youth in New York averaged \$1,874 per underage customer. Underage customers were heavier consumers than adults. They drank an average of 4.0 drinks per day; in contrast, legal customers consumed only 1.5.

Harm Associated with Underage Drinking in New York

Underage drinking in New York leads to substantial harm due to traffic crashes, violent crime, property crime, unintentional injury, and high-risk sex.

- During 2012, an estimated 46 traffic fatalities and 1,873 nonfatal traffic injuries were attributable to driving after underage drinking.
- In 2012, an estimated 44 homicides; 31,600 nonfatal violent crimes such as rape, robbery, and assault; 33,000 property crimes including burglary, larceny, and car theft; and 618,000 public order crimes including vandalism, disorderly conduct, loitering, and curfew violations were attributable to underage drinking.
- In 2011, an estimated 12 alcohol-involved fatal burns, drownings, and suicides were attributable to underage drinking.
- In 2013, an estimated 1,152 teen pregnancies and 46,563 teens having high-risk sex were attributable to underage drinking.

For comparison with other states, in U.S. rather than state prices, the harm from underage drinking per youth in New York averages \$861. Such comparisons require caution. In part, they may reflect differences in crime and crash rates, problem-reporting to police, and co-occurring drug use.

Produced by the Pacific Institute for Research and Evaluation (PIRE) with funding from the Office of Juvenile Justice and Delinquency Prevention (OJJDP), March 2015.

¹ Taylor DM, Miller TR. (2015). Methodology: Underage Drinking Fact Sheets. Calverton, MD: PIRE, <http://www.udetc.org/documents/Underage-Cost-Methods-082807.pdf>

² Office of Applied Studies, Substance Abuse and Mental Health Services Administration. Treatment Episode Data Set. (2013). *Substance Abuse Treatment by Primary Substance of Abuse, According to Sex, Age, Race, and Ethnicity, 2011*. Available [Online]: <http://www.icpsr.umich.edu/icpsrweb/SAMHDA/studies/30462>

³ Grant, B.F., & Dawson, D.A. (1997). Age at onset of alcohol use and its association with DSM-IV alcohol abuse and dependence: Results from the National Longitudinal Alcohol Epidemiologic Survey. *Journal of Substance Abuse* 9: 103-110.

⁴ Centers for Disease Control (CDC). (2013). Youth Risk Behavior Surveillance System (YRBSS). Available [Online]: <http://nccd.cdc.gov/youthonline/App/Default.aspx>. Or an equivalent state data system.

⁵ Lifetime use was based on imputation using YRBS state reports.